



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

of a sanitary inspector, resident in a provincial capital, whose duties shall be to enforce all laws and ordinances of public health and to carry out the instructions of the director general. A sanitary inspector must be a graduate of medicine and surgery, and must have had at least three years of practice.

DEATHS DURING WEEK ENDED APRIL 5, 1919, IN CITIES.

The following table shows the registered deaths from all causes and from pneumonia (all forms) and influenza combined, in certain large cities of the United States during the week ended April 5, 1919. The annual death rates per 1,000 population for the week and for the corresponding week of previous years are also shown. Weekly figures fluctuate widely and caution must be used in their interpretation.

The data are taken from the "Weekly Health Index," April 8, 1919, issued by the Bureau of the Census, Department of Commerce. The populations used in computing the rates are estimated by the Bureau of the Census as of July 1, 1918.

Registered deaths and annual death rates per 1,000 population in certain large cities of the United States, week ended April 5, 1919—Deaths from all causes and from pneumonia (all forms) and influenza combined.

City.	Population July 1, 1918, estimated.	Total deaths, all causes.	Annual death rate per 1,000.	Annual death rate for preceding years. ¹	Influenza and pneumonia (all forms).	
					Number of deaths.	Annual death rate per 1,000.
Albany, N. Y.	112,565	53	24.6	C. 21.3	19	8.8
Atlanta, Ga.	201,732	58	15.0	O. 15.8		
Baltimore, Md.	² 669,981	211	16.4	A. 20.3	28	2.2
Birmingham, Ala.	197,670	61	16.1	A. 18.0		
Boston, Mass.	785,245	240	15.9	A. 17.8	33	2.2
Buffalo, N. Y.	473,229	163	18.0	C. 19.4	28	3.1
Cambridge, Mass.	111,432	32	15.0	A. 16.5	5	2.3
Chicago, Ill.	2,596,681	790	15.9	A. 16.4	170	3.4
Cincinnati, Ohio.	418,022	147	18.3	C. 24.8	40	5.0
Cleveland, Ohio.	810,306	218	14.0	C. 13.6	71	4.6
Columbus, Ohio.	225,296	83	19.2	C. 21.8	29	6.7
Dayton, Ohio.	130,655	41	16.4	C. 25.9		
Denver, Colo.		78		A.	10	
Fall River, Mass.	128,392	25	10.2	C. 14.2	3	1.2
Grand Rapids, Mich.	135,450	35	13.5	C. 13.9		
Indianapolis, Ind.	289,577	109	19.6	C. 16.9		
Jersey City, N. J.	318,770	97	15.9	C. 21.6		
Kansas City, Mo.	313,785	107	17.8	C. 24.1	36	6.0
Los Angeles, Calif.	568,495	168	15.4	A. 12.8	19	1.7
Louisville, Ky.	242,707	71	15.3	C. 22.3	16	3.4
Lowell, Mass.	109,081	37	17.7	A. 17.7	6	2.9
Memphis, Tenn.	154,759	79	26.6	C. 20.9	12	4.0
Milwaukee, Wis.	453,481	135	15.5	A. 13.8	48	5.5
Minneapolis, Minn.	383,442	85	11.6	C. 21.2		
Nashville, Tenn.	119,215	49	21.4	C. 20.1	12	5.2
New Haven, Conn.	154,865	45	15.2	C. 17.8		
New Orleans, La.	382,273	131	17.9	A. 21.2		
New York, N. Y.	5,215,879	1,774	17.7	C. 21.6	480	4.8
Oakland, Calif.	214,206	40	9.7	A. 11.1		
Philadelphia, Pa.	1,761,371	582	17.2	³ 18.8	98	2.9
Pittsburgh, Pa.	593,303	206	18.1	C. 24.3	75	6.6
Portland, Oreg.		58		C.	11	
Providence, R. I.	263,613	83	16.4	C. 21.8	14	2.8
Richmond, Va.	160,719	45	14.6	C. 19.5	7	2.3
Rochester, N. Y.	264,856	79	15.6	C. 16.1		
St. Louis, Mo.	779,951	201	13.4	C. 15.3	54	3.6
St. Paul, Minn.	257,699	69	14.0	C. 15.8		
San Francisco, Calif.	478,530	142	15.5	C. 15.5	27	2.9
Seattle, Wash.		67		A.	8	
Spokane, Wash.		29		C.	4	
Syracuse, N. Y.	161,404	52	16.8	C. 24.9	6	1.9
Toledo, Ohio.	262,234	76	15.1	A. 16.2	36	7.2
Washington, D. C.	401,681	131	17.0	A. 18.6	17	2.2
Worcester, Mass.	173,650	52	15.6	C. 24.9		

¹ "A" indicates that the rate given is the average annual death rate per 1,000 population for the corresponding week of the years 1913 to 1917, inclusive. "C" indicates that the rate is the annual death rate per 1,000 population for the corresponding week of 1918.

² Population estimated as of July 1, 1919.

³ Rate is based on statistics of 1915, 1916, and 1917.